

Fisher success and adaptation to plantation systems in Chile

Tracy Van Holt

East Carolina University

**Department of Geography & Institute for
Coastal Science & Policy**

September 26, 2012

vanholtt@ecu.edu

Chilean Fishers 30 organizations

Carlos Moreno, Universidad Austral

Sandor Mulsow, Universidad Austral

Michael Binford, University of Florida

H. Russel Bernard, University of Florida

Tom Frazer, University of Florida

Kenneth Portier, American Cancer Society

Rodrigo Vergara, University of Florida

Financial Support:

NASA Earth System Science Fellow

National Security Education Program

Rotary Ambassadorial Scholarship

Chilean Fisheries Management



Loco (*Concholepas concholepas*)


Chilean Fisheries Management

Sustainable Harvest	Crash	Recovery (Management Areas)
1970's	1980's	2000's



Consequences of Territorial User Rights Fisheries

1. Locos available for harvest
2. Restricted access & movement
3. Introduced new fishers (knowledge)
4. Loco biological condition & price varies



Territorial User Rights Fisheries MA³

Skipper Effect Theory



- Some fishers catch more fish than others
- Explained by experience and/or technology
- Management implications overfishing



1-What factors account for fisher success?

2-How are fishers adapting to environmental change?



The fisheries

Loco
(*Concholepas concholepas*)




closed access

Congrio
(*Genypterus* sp.)



open access

Research Design: space for time substitution




southern Chile • similar coastal shelf • little el niño influence & upwelling • similar benthic & fish fauna • landscape change varies

(Fernandez et al. 2000; Lancellotti & Vasquez 2000; Camus 2001)


Forest Plantations Increased from 1985-2001

Plantaciones forestales se incrementaron desde 1985 al 2001




Supervised classification of October 5, 1985 (Landsat TM, path 233, rows 87-89) and November 29, 2001 (Landsat ETM, path 233, rows 87-89). Images were atmospherically and topographically corrected. The numbers in white boxes correspond to watersheds in study site.

Chlorophyll-a concentration higher in nearshore influenced by plantations




SeaWiFS satellite images from winter (April to July) 2003/Imágenes satelitales SeaWiFS del invierno (Abril-Julio) del 2003


Loco shells from plantation-influenced watersheds have more epibionts & endobionts.



Phoronids



Barnacles



Polychaetes



Bivalves



clean shell

Northern Study Site

